

Exacerbation of COPD and what to expect after discharge

Introduction

This leaflet gives you information about COPD and how you can help to reduce the chances of another attack.

What is COPD?

COPD (Chronic Obstructive Pulmonary Disease) is a group of chronic lung conditions that cause permanent narrowing of the airways and difficulties with taking in oxygen and removing carbon dioxide from the blood stream. COPD is usually the result of inhalation of harmful substances, most commonly cigarette smoke. There are some rare genetic conditions that may cause COPD, mainly in younger patients.

The most important thing to do if you have COPD is to stop smoking. If you are a current smoker and are keen to stop, there are details how to access support at the end of this leaflet.

Exacerbation of COPD

For many people their COPD is manageable on a day to day basis. However, from time to time, things can temporarily get worse – this is an exacerbation. Exacerbations can be triggered by a number of factors such as infection, changes in the weather or exercise.

Often exacerbations can be managed at home with increased use of inhalers and a short course of steroids.

If very severe, COPD exacerbations can be life-threatening and need urgent treatment in hospital. Treatment may include oxygen and nebulisers.

If the cause of the exacerbation is thought to be an infection, a course of antibiotics may be needed.

Reference No.

GHPI1644_02_21

Department

Respiratory

Review due

February 2024

Patient Information

More information about COPD can be found at:

British Lung Foundation

Website: www.blf.org.uk/support-for-you/copd

This website has useful advice regarding a wide range of issues related to COPD.

What to do after an exacerbation of COPD?

Your symptoms should start to improve 2 to 3 days after starting treatment.

You should complete any course of steroids or antibiotics you have been prescribed. If you are taking long term steroids (usually prednisolone) or have been on a prolonged course, the dose may need to be slowly reduced rather than stopping abruptly.

While your breathing should get better after an exacerbation, it often does not quite get back to the same level as before.

Signs that might suggest things are getting worse or you are developing another flare up are:

- Increased breathlessness that does not get better after a short time
- Increased coughing
- Producing more sputum
- Your sputum changes in consistency or colour.

Patient Information

Do I feel worse than usual?

If I get:



more out of breath



more out of breath despite taking my reliever medicines



more phlegm, change in colour: dark yellow, green



more chesty coughing



I will:



use my reliever inhaler more



start my rescue pack - steroids



start my rescue pack - steroids and antibiotics



keep calm and use ways to control my breathing

I will remember:

to tell my doctor or nurse



that if I feel really bad, I must call 999



If I use my rescue pack, I will tell my doctor or nurse so they can keep track of my flare-ups and I can get my medicines replaced.

Follow up arrangements

Your inhaler treatment and need for steroids and antibiotics should have been explained before you are discharged from hospital. If you did not understand the information or you have some concerns or questions, please ask your nurse or doctor before leaving.

Patient Information

It is recommended that you should have a review of your COPD in the first 4 weeks after discharge. This would usually be with your GP or COPD nurse. The purpose of this review is to check your symptoms and make sure that you are on the correct medication. Your GP or COPD nurse will also check that you have been shown the best way to use your inhalers.

A further review is recommended at 12 to 16 weeks. This is similar to the above review, but should also provide a good opportunity for your lung function to be assessed with some breathing tests (called spirometry). This will be arranged through your GP, but may take place at a community health centre rather than at the practice.

If a hospital follow up appointment is needed it will be arranged and the details sent to you.

Smoking cessation support

If you are a current smoker and are keen to give up, support is available from Healthy Lifestyles. You can access this support either through the website or by telephone.

Website: www.hlsglos.org

Tel: 0800 122 3788

Content reviewed: February 2021