

Patient
 Information

Gastroscopy with Radiofrequency Ablation (RFA) for oesophageal dysplasia

Information

You have been offered a treatment to your oesophagus (gullet) called Radio Frequency Ablation (RFA). This leaflet contains information about the treatment and will hopefully answer any questions that you may have. It is important that you read this information so that you understand why you have been recommended for this treatment and what it involves.

If you are pregnant, taking warfarin, other blood thinning drugs or any medication for diabetes, please contact the Medication Advice Line as soon as possible for advice. The number is at the end of this leaflet. Your test may be delayed if you do not do so.

What is dysplasia in the oesophagus?




Figure 1: GI tract

Reference No.

GHP11215_03_20

Department

Endoscopy

Review due

March 2023

Patient Information

Dysplasia is the term used to describe cells in the lining of the oesophagus that look abnormal under a microscope. It is common in patients with 'Barrett's Oesophagus'. The severe forms of dysplasia (called 'high grade dysplasia') can signify that the cells have the potential to become cancerous. Oesophageal cancer is a serious condition, so we try to catch any changes at an early stage when it can be successfully treated.

What is the aim of RFA?

Dysplasia and very early oesophageal cancer affects only the cells lining the oesophagus.

When we perform the RFA, we can destroy these abnormal cells; once performed we expect the oesophagus lining to heal with normal cells. By treating the abnormal cells and preventing the development of cancer we would hope to prevent the need for major surgery.

Preparing for your procedure

To allow a clear view during the procedure your stomach must be empty. You will be asked not to eat anything for at least 6 hours before your appointment time. You may drink water up to 2 hours before the appointment time.

On admission to the Endoscopy Unit

- You will be seen by a nurse who will check your personal details
- You will be asked a series of questions including any operations or illnesses that you have had or are presently suffering with
- Please bring a list of your medications with you
- The nurse will want to know if you have any allergies or bad reactions to any medications
- This procedure will be carried out with sedation
- You will need a responsible adult to take you home when you are ready for discharge
- You should also have a responsible adult with you at home for 24 hours after your procedure

Patient Information

During the procedure

In the treatment room you will have a local anaesthetic spray applied to the back of your throat. You will be asked to remove any glasses and/or dentures and monitoring equipment will be attached.

You will then lie on your left hand side. A plastic tube (also called a cannula) will be placed into a vein in your hand or arm. This will be used to give you the sedative drugs. As a safety measure you will be given oxygen via a nasal sponge. • Once you are feeling sleepy, the endoscope is passed down into your oesophagus to look carefully at the area that needs to be treated

- Once the area has been assessed and we have confirmed it is suitable for the procedure, a guiding wire is passed through the endoscope into your stomach
- The endoscope is then taken out of your oesophagus and a balloon is passed down over the guiding wire into your oesophagus. This allows us to select the correct size of balloon required to deliver the radiofrequency energy
- The first balloon is removed and another balloon for delivering the treatment is passed over the guiding wire

The procedure will take about 45 minutes, but this does depend on the size of area we are treating.

Complications and risks

RFA has been used to treat patients from all over the world for several years. It has been shown to be a very safe procedure.

As with all endoscopy techniques, there are small risks involved. The major risks can include damage to the oesophagus (perforation) and causing bleeding from the oesophagus. The risk of this happening is about 1 in 1000.

After the procedure

After we have completed your procedure you will be taken, on your trolley, to the recovery area. You will continue to be given some oxygen, and your blood pressure will be monitored until you are awake.

Patient Information

You should not experience any significant discomfort after the procedure but you might be aware of a feeling like heartburn or indigestion. You will be given a drink when you are awake.

On admission

On admission the nursing staff will ask you some questions and make sure that you are prepared for your procedure. You will also have the opportunity to ask any questions you may have about what is going to happen.

Going home

You will have been given a sedative for the procedure, so you must arrange for someone to take you home. You will also need a responsible adult to stay with you for 24 hours. For this period you should not:

- drive a car, motorbike or ride a bicycle
- drink alcohol
- operate machinery or do anything requiring skill or judgement, including cooking
- make important decisions or sign important documents

What can I eat after the procedure?

For the rest of the day following the procedure, you should drink fluids only. Do not drink anything too hot or cold, room temperature is about right. The day after the procedure, you should start on a soft, sloppy diet and stay on this for a week. This includes breakfast cereals and other food you do not have to chew too much such as mashed potato, minced meat and pasta.

Medication

You should carry on taking your medicine for reducing stomach acid as usual, but for the week after the procedure you should take double your normal amount (dose). You can also take Gaviscon® (this should be Gaviscon Advanced®) or other antacid liquids, if required.

Patient Information

If you experience mild chest discomfort you can take pain relief such as soluble paracetamol.

Emergency contact

If you develop any of the following symptoms, you should contact your GP, or if out of normal surgery hours NHS 111 immediately.

- Severe chest pain
- Shortness of breath
- High temperature
- Vomiting or vomiting blood
- Abdominal distension

Follow up

You will receive an appointment in the post to return for another endoscopy to allow us to check that the RFA has been successful. This is usually 2 to 3 months after your procedure. If further treatment is needed, this will be given during this later endoscopy.

Contact information

If you have any questions or concerns about your appointment date or time, please contact the Booking Office.

Booking Office

Tel: 0300 422 6350

Monday to Friday, 8:00am to 4:00pm

Appointment enquiries booking team

Cheltenham General Hospital

Tel: 0300 422 6899

Monday to Friday, 8:30am to 4:00pm

Gloucestershire Royal Hospital

Tel: 0300 422 6351

Monday to Friday, 8:30am to 4:00pm

**Patient
Information**

Medication Advice Line (answer machine)

If you have any questions relating to your medication, please leave a message and a member of staff will return your call:

Cheltenham General Hospital

Tel: 0300 422 3370

Gloucestershire Royal Hospital

Tel: 0300 422 8232

Other Endoscopy Units

Cirencester Hospital

Tel: 0300 422 6294

Stroud General Hospital

Tel: 0300 421 8073

NHS 111

Tel: 111

Content reviewed: March 2020