

**Patient
Information**

Pulled elbow

Introduction

You have been given this leaflet because your child has been examined and diagnosed with a pulled elbow. This type of injury may also be called a radial head subluxation, nursemaid elbow or annular ligament displacement

What is a pulled elbow?

Pulled elbow is a common injury suffered by children usually between the age of 1 and 5 years. Young children's ligaments (the elastic-like bands that hold bones together) are slightly loose so it is easy for a ligament in the elbow to slip into the joint and get stuck.

It is unusual for children over 5 years old to get a pulled elbow, as their ligaments are less stretchy. In most cases, children with a pulled elbow will cry straightaway and will not use the injured arm at all, or they will simply have it hanging by their side.

What causes a pulled elbow?

It is often the result of a sudden jerking or pulling action on the child's arm or forearm, for example:

- Pulling a child up by the hands
- Swinging a toddler
- Jerking an arm when pulling a toddler

This sudden tug on the arm pulls the radius through the annular ligament which may partially tear and become trapped between the radial head and the capitellum.

He or she will keep the arm in a straight position or with a slight bend in the elbow. The injury will not be obvious because a pulled elbow does not usually cause arm shape changes or swelling.

This is not the same as a dislocation where the bones in the joint are not lined up correctly.

Reference No.

GHPI0839_01_19

Department



**Emergency
Department**

Review due

January 2022

**Patient
Information**

Diagram of a child's elbow joint



**Patient
Information**

Treatment

The arm will be moved or manipulated (put back into place) by a nurse or doctor; this will only take a few seconds. The child will be able to sit on a parent or carer's knee for the procedure. The doctor will then gently take the arm from a straight position and bend it upwards or straighten the arm while turning the palm to the floor.

The child will usually feel a brief moment of pain during the procedure, but quickly feel much better. Most have full use of the arm within 5 to 10 minutes. Some children may need this procedure to be repeated (especially if the pulled elbow happened less recently) before the ligament returns to normal.

Often children may not want to use the arm after the procedure, fearing it will be painful but once they have their attention on playing they will forget the experience.

Will my child need an X-ray?

If the nurse or doctor is sure that the injury is a pulled elbow, an X-ray is not needed as there is no injury to the bones.

Following discharge

Usually your child will not need pain relief medication but if the elbow was 'out' for a while, then your child may need some pain relief medication on the day of the treatment. If your child is not moving their arm fully by the next day, they will need to return either to a Minor Injuries Unit or Emergency Department.

Children who have had a pulled elbow might get it again. So be mindful of the risks and do not pull, tug, or swing your child by the arms or hands.

Please give this advice to anyone who provides cares for your child.

Please visit the NHS Choices website for further information.

**Patient
Information**

Further information

Further information may be obtained via your GP, NHS 111 or NHS choices.

NHS 111

Tel: 111

NHS Choices

Website:

www.nhs.uk/conditions/arpain/Pages/Introduction.aspx

Content reviewed: January 2019

Patient
Information

**Patient
Information**

**Patient
Information**

**Patient
Information**