

**Patient
Information**

Surgical insertion of a Peritoneal Dialysis (PD) catheter

Introduction

You have been given this leaflet because your kidneys are no longer functioning efficiently. Excess water and waste needs to be removed from your body in another way, through dialysis. The method of dialysis you have chosen is peritoneal dialysis. To enable you to have this type of dialysis, you need a Peritoneal Dialysis (PD) catheter.

What is a Peritoneal Dialysis (PD) catheter?

The PD catheter is a soft silicone tube placed inside the abdomen. This allows dialysis fluid to enter the abdominal cavity and then drain back out again.

What are the benefits of having the PD catheter inserted?

The PD catheter will be inserted while you are under a general anaesthetic (while you are asleep).

If you have had previous operations on your tummy, it may be too uncomfortable to have the procedure without a general anaesthetic.

A general anaesthetic is also preferred if you are having another procedure done at the same time (for example a hernia repair).

Pre-Assessment Clinic

You will need to visit the Pre-Assessment Clinic and be examined by the surgeon or a member of the surgical team before a date for the procedure can be arranged. The surgeon will discuss the procedure with you and answer any questions you may have.

Reference No.

GHP11345_10_20

Department

Renal Services

Review due

October 2023

Patient Information

Before any hospital admission, and to reduce the risk of infection, you will be screened for Methicillin-Resistant Staphylococcus Aureus (MRSA). You will be given an antiseptic skin wash. This is to be used for the 5 days before your operation to reduce any bacteria on your skin.

Where will the PD catheter insertion be carried out?

The procedure will be carried out in an operating theatre either at Gloucestershire Royal Hospital or Cheltenham General Hospital.

How long will the procedure take?

The procedure will take about 45 minutes, but you also need to add the time it takes to be anaesthetised and recover.

Do I need to prepare for the procedure?

You will be asked to shower or bath at home, using the antiseptic wash for 5 days, before the day of the procedure.

You will also need to take a strong laxative as advised before the procedure; this is to make the bowels loose.

You must not eat or drink for 6 to 8 hours before having a general anaesthetic. Do not suck sweets or chew gum during this period. Full instructions will be given in the letter you will receive confirming your operation date and time.

Valuables

Please do not bring valuables, jewellery or large sums of money into the hospital. If this is unavoidable, please ask a relative or a friend to take them home for you. The hospital cannot accept responsibility for the loss of items.

What happens after the procedure?

You may need to stay overnight if the procedure is not carried out until the afternoon. You will then be discharged the next morning. If you have the operation in the morning you may be able to go home early evening if there are no complications.

Patient Information

Do not travel home by public transport after you have had your PD catheter inserted. You should ask a relative or friend to drive you home by car. You should not drive yourself for at least for 2 weeks. If you cannot arrange your own transport please let us know, before the day of your appointment, so that we can book transport for you.

Will I have any stitches?

Surgical glue or stitches will be used to close your wound. We will advise you of the type of wound closure used. If you have had stitches then we will arrange for them to be removed when needed.

Are there any risks?

There are very few risks associated with having a PD catheter inserted, but you should be aware of the possible complications.

When the PD catheter is being inserted, it may accidentally puncture the wall of the bowel or bladder. If this happens, the catheter is removed immediately but you might need to have an additional operation to repair your bowel. This is a very rare complication.

After it has been inserted, your PD catheter may move into a poor drainage position, especially if your bowels are not loose (remember to take your laxative as directed). Your tube may need to be repositioned. This complication happens in 10 to 25 of every 100 cases.

The fluid we use to flush the catheter after it has been inserted may appear blood stained. This usually stops without further treatment.

You may develop an infection. We will tell you how to recognise the signs of an infection before you are discharged. If you do develop an infection, you will need to take antibiotics. Rarely, you may need to have your catheter taken out and another one inserted a few weeks later.

There is always a small risk of a serious reaction or complication if you have a general anaesthetic.

**Patient
Information**

What should I do if I cannot attend my appointment or I am not feeling well?

Please let us know as soon as you can if you are unable to attend your appointment, you are experiencing diarrhoea/vomiting or have a high temperature so we can advise you and offer another appointment.

Contact information

If you have any questions or queries, please contact:

PD Specialist Nurses

Tel: 0300 422 6761

Monday to Friday, 8:00am to 4:00pm

Ward 7b, Renal Unit

Gloucestershire Royal Hospital

Tel: 0300 422 6768

Available 24 hours a day

PD Coordinator

Tel: 0300 422 6890

Monday to Friday, 7:30am to 4:30pm

Mr Cooper's secretary

Tel: 0300 422 3345

Monday to Friday, 8:00am to 4:00pm

Mr Kulkarni's secretary

Tel: 0300 422 3355

Monday to Friday, 8:00am to 4:00pm

Mr Paravastu's secretary

Tel: 0300 422 3331

Monday to Friday, 8:00am to 4:00pm

Further information

Kidney Patient Guide

Website: www.kidneypatientguide.org.uk

Content reviewed: October 2020